Hydrobiologia **355:** v–vi, 1997.
A.D. Naumov, H. Hummel, A.A. Sukhotin & J.S. Ryland (eds),
Interactions and Adaptation Strategies of Marine Organisms.

Contents

Preface vii

Interactions in Marine Organisms

The importance of juveniles in structuring a littoral macrobenthic community
by I.V. Burkovsky, A.A.Udalov & A.P. Stoljarov 1–9

Epibenthic predation in marine soft-bottoms: being small and how to get away with it
by M. Thiel 11–19

Abundance, feeding behaviour and nematocysts of scyphopolyps (Cnidaria) and nematocysts in their predator, the nudibranch *Coryphella verrucosa* (Mollusca)
by C. Östman 21–28

The importance of intraspecific competition in a *Littorina littorea* population in the Wadden Sea
by C. Fenske 29–39

Occurrence of epifauna on the periwinkle, *Littorina littorea* (L.), and interactions with the polychaete *Polydora ciliata* (Johnston)
by G.F. Warner 41–47

Effects of epibiosis on consumer–prey interactions
by M. Wahl, M.E. Hay & P. Enderlein 49–59

Parasites on an intertidal *Corophium*-bed: factors determining the phenology of microphallid trematodes in the intermediate host populations of the mud-snail *Hydrobia ulvae* and the amphipod *Corophium volutator*
by K.N. Mouritsen, T. Jensen & K.T. Jensen 61–70

The association between the caprellid *Pariambus typicus* Krøyer (Crustacea, Amphipoda) and ophiuroids
by U. Volbehr & E. Rachor 71–76

Chemically-mediated interactions in benthic organisms: the chemical ecology of *Crambe crambe* (Porifera, Poecilosclerida)
by M.A. Becerro, M.J. Uriz & X. Turon 77–89

Fauna associated with detached kelp in different types of subtidal habitats of the White Sea
by A.B. Tzetlin, V.O. Mokievsky, A.N. Melnikov, M.V. Saphonov, T.G. Simdyanov & I.E. Ivanov 91–100

Soft-bottom macro invertebrate fauna of North Norwegian coastal waters with particular reference to sill-basins. Part one: Bottom topography and species diversity
by L.-H. Larsen 101–113

Adaptation Strategies in Marine Organisms

Mechanisms of salinity adaptations in marine molluscs
by V.J. Berger & A.D. Kharazova 115–126
Sensitivity to stress in the bivalve *Macoma balthica* from the most northern (Arctic) to the most southern (French) populations: low sensitivity in Arctic populations because of genetic adaptations?
by H. Hummel, R. Bogaards, T. Bek, L. Polishchuk, C. Amiard-Triquet, G. Bachelet, M. Desprez, P. Strelkov, A. Sukhotin, A. Naumov, S. Dahle, S. Denisenko, M. Gantsevich, K. Sokolov & L. de Wolf 127–138
Defenses against oxidative stress in the Antarctic scallop *Adamussium colbecki* and effects of acute exposure to metals
by F. Regoli, M. Nigro, E. Bertoli, G. Principato & E. Orlando 139–144
A new species of *Hyalopomatus* (Serpulidae: Polychaeta) which lacks an operculum: is this an adaptation to low oxygen?
by E.W. Knight-Jones, P. Knight-Jones, P.G. Oliver & A.S.Y. Mackie 145–151
Adaptation capabilities of marine modular organisms
by N.N. Marfenin 153–158
Morphological convergence of resting stages of planktonic organisms: a review
by G. Belmonte, A. Miglietta, F. Rubino & F. Boero 159–165
Effects of experimental conditions on the feeding rate of *Mysis mixta* (Crustacea, Mysidacea)
by E. Gorokhova & S. Hansson 167–172
Feeding behaviour of *Cerastoderma edule* in a turbid environment: physiological adaptations and derived benefit
by M.B. Urrutia, J.I.P. Iglesias & E. Navarro 173–180